

Lentil Variety Development for Yield and Disease Resistance for Potential Areas: Registration of a Lentil Variety Named Debine

Amanuel Tekalign*, Tadele Tadesse, Belay Asmare

Oromia Agriculture Research Institute, Sinana Agriculture Research Center, Bale-Robe, Ethiopia

Email address:

amnu2012@gmail.com (A. Tekalign)

*Corresponding author

To cite this article:

Amanuel Tekalign, Tadele Tadesse, Belay Asmare. Lentil Variety Development for Yield and Disease Resistance for Potential Areas:

Registration of a Lentil Variety Named Debine. *Computational Biology and Bioinformatics*. Vol. 10, No. 1, 2022, pp. 9-13.

doi: 10.11648/j.cbb.20221001.12

Received: February 22, 2022; **Accepted:** March 28, 2022; **Published:** March 31, 2022

Abstract: The development of new varieties with high yield and acceptable levels of stability is an important in breeding program. The performance of a given genotype depends on its genetic potential and the environment upon which it is grown. Debine is a commercial name given for a newly released Lentil (*Lens culinaris*) variety with pedigree designation of ‘DZ - 2012-LN-0051’ was released in 2021 for highland areas of Bale, Southeastern Ethiopia and other similar agro ecologies. The variety has been registered by Sinana Agricultural research center and it was tested at better representative environments (Sinana and Agarfa) representing highland (2300 to 2600) meter above sea level) agro-ecologies during 2016 to 2018 main cropping season. The variety is mainly characterized by its superior mean grain yield as compared from standard check Asano based on different yield measurement and stability testing parameters across locations and over years. It also had comparable resistance/tolerant level to major Lentil diseases such as Aschocyta blight, Rust and Root Rot. Debine has dark gray seed color and could be cultivated across a number of locations in the highlands of Bale and other similar agro-ecologies for increasing productivity of the crop.

Keywords: Disease Resistance, Grain Yield, Lentil (*Lens culinaris*), Stability, Variety Registration

1. Introduction

Lentil (*Lens Culinaris* Medikus.; Fabaceae) is an autogamous, diploid ($2n = 2x = 14$), self-pollinated and annual cool season grain legume with ~4 Gbp genome size [10, 2]. It is one of the first agriculture crop grown more than 8500 years ago [1, 3]. Its annual production is ~5 million ton globally and Canada, India, Turkey, USA, Nepal, Australia, Ethiopia, Bangladesh, Kazakhstan is the major lentil producing countries of the world sequentially. Approximately 50% of world’s lentils are grown in South Asia, and nearly 1.5 billion people in this area consume ~70% of the global lentil supply [2, 5].

Lentil plays a significant role in human and animal nutrition and in maintenance and improvement of soil fertility [8, 9]. Its cultivation enriches soil nutrient status by adding nitrogen, carbon and organic matter which promotes

sustainable cereal-based systems of crop production [8]. It is a nutritious food legume. It is cultivated for its seed and mostly eaten as split [6]. The primary product of lentil is its seed which has relatively higher contents of protein, carbohydrate and calories compared to other legumes [7]. It is the most desired crop because of its high average protein content and fast cooking characteristic in many lentil producing regions. It can be used as a main dish, side dish, or in salads. Seeds can be fried and seasoned for consumption but sometimes difficult to cook because of the hard seed coat those results from excessive drying [12]. Its flour is used to make soups, stews, purees, and mixed with cereals to make bread and cakes; and as a food for infants [11].

Because of its significant economic role and social conditions, lentil production has recently been expanding in

both stressed and non-stressed environments. In Ethiopia in 2017 cropping season, the total area of production of pulse crops is about 2,092,357.57 hectares and the total production was 328,734.78ton. Among these pulse crops, lentil covered 124,915.16 hectares with production of 170.09ton. The national productivity of lentil was 1.44 t ha⁻¹ [4], which was far below the potential yield of the crops and productivity in different parts of the world. The reason for this yield gap is mainly due to poor genetic makeup of the available cultivars, and other biotic and abiotic factors. Therefore, the objective of this study was to register stable high yielding and disease

resistant/tolerant Lentil variety for highlands Bale and other similar agro-ecologies in Ethiopia.

2. Materials and Methods

2.1. Descriptions of Experimental Sites

The experiment was conducted at two potential areas of Bale Zone, Sinana and Agarfa in South Eastern Ethiopia. Description of the study sites at Regional Variety Trail is given below.

Table 1. Description of the test locations for geographical position and physico-chemical properties.

Parameter	Location	
	Sinana	Agarfa
Geographical position		
Latitude	07°07' N	07°15'44''N
Longitude	40°10'00''E	039°50'38''E
Altitude (m.a.s.l.)	2400	2509
Soil property		
pH	6.2	6.3
Texture	Clay	Clay loam
OMC (%)	3.9	3.4
Total N (%)	0.24	0.2
Pav (ppm)	30.4	32.41
K (mg/Kg)	240	572
CEC (meq/Kg)	64.4	71.5
Moisture Regime	SH2	SH2

Key: OMC = Organic matter content, N = Nitrogen, Pav = Phosphorus availability, K = Potassium, CEC = Cation exchange capacity, SH2 = Sub-humid tepid to cool sub-humid mid-highlands.

2.2. Breeding Procedures

Adapted Lentil line “Dz-2012-Ln-0051, which was selected from the last stage of variety trial. The crossing was done at Debre Zeit Agricultural Research Center. Screen houses were routinely used in the early generations, i.e., F1, F2, F3 and F4, of a breeding cycle. During these phases, selection for traits with high heritability such as; seed size, grain yielding ability, plant habit, time of flowering and resistance to major diseases were undertaken. selected individual lines from F5 generation were evaluated for yielding ability, large seed size, disease reaction and stability in preliminary yield trial (PYT) conducted at Sinana Agricultural Research Center. From this trial, 15 promising genotypes were promoted and evaluated in a regional variety trial (RVT) along with standard checks ‘Asano’ at multi-locations (Sinana and Agarfa) during 2016 to 2018 main cropping seasons. Lastly, Dz-2012-Ln-0051 and Dz-2012-Ln-0085 were selected as the most promising candidate varieties and verified along with best standard checks ‘Asano’ on 10 m x 10 m plots. National Variety Release Technical Committee were evaluated each one on-station and two on-farm fields during the 2020/21 cropping season. Finally, the committee decided the first genotypes coded as Dz-2012-Ln-0051, and named “Debine”, for official release.

3. Result and Discussions

3.1. Agronomic and Morphological Characteristics

In an attempt to develop Debine, higher yield, and resistance to major lentil diseases were important traits of consideration. The newly released Lentil variety ‘Debine’ is characterized by an erect growth habit. Its flower color is light Pink. The seed coat and cotyledon colors are dark gray and light red, respectively. The average number of days required to reach its 50% flowering and 95% physiological maturity were 62 and 122, respectively, with the average plant height being 32 cm (Table 2). The average number of pods per plant is 33 (Table 4). It has good general acceptance for lentil with high quality. The appropriate planting date for this variety would range from end of July to early August (Table 2).

3.2. Yield and Quality Performance

Highly significant variations among Lentil genotypes were observed throughout the trial evaluation. Debine consistently out-yielded other tested Lentil genotype over three years. Combined location over years analysis revealed that it had produced an average yield of 22-25 Q/ha at Research field and 14-16 Q/ha on farm yield. This means that the grain yields of Debine was found to be 29.76% yield advantage

over standard check Asano (Table 4). Debine offers new hope for resource poor farmers in study areas and other similar agro ecology.

3.3. Reaction to Major Diseases

Developing Lentil cultivars with high yielder, resistant or tolerant varieties to major lentil diseases such as *Ascochyta lentis* (*ascochyta blight*), Rust (*Uromyces viciae-fabae*) and Root Rot is among the major objectives of the Lentil breeding program. Accordingly, above mentioned disease is among the major bottleneck for Lentil production in Southeastern part of the country, Bale. Disease data across location and years were scored and analyzed. Debine variety showed resistance to moderate resistance to the above-mentioned diseases throughout the field evaluation periods (Table 5).

3.4. Performance Stability and Adaptation Domain

The variety 'Debine' was released for high altitude agro-ecologies of the country receiving 750-to-1000 mm average annual rainfall. It is well adapted to an altitude range of 1800 – 2600 meters above sea level such as Sinana, Goba, Agarfa, Gassera, Goro (Meliyu), Adaba, Dodola and other similar agro-ecologies (Table 2). Based on most stability parameters, 'Debine' showed relatively comparable performance stability across a range of environments (Table 4).

3.5. Variety Maintenance

The breeder and foundation seed will be maintained by Sinana Agricultural Research Center/ Oromia Agricultural Research Institute.

Table 2. Agronomical and Morphological Characteristics and Agro-ecological Zones of Adaptation of Debine, Lentil variety.

No	Variety name:	Debine (DZ-2012-LN-0051)
1	Adaptation area	Sinana, Goba, Agarfa, Gassera, Goro (Meliyu), Adaba, Dodola and other similar agro-ecologies
2	Altitude (m.a.s.l.)	1800 – 2600
3	Rainfall (mm)	750 – 1000
4	Seed Rate (Kg/ha)	65
5	Planting date	End of July to Early August
6	Days to Flower	62
7	Days to Maturity	122
8	Plant Height (cm)	32
9	Growth habit	Erect
10	1000 Seed Weight (gm)	3.7
11	Seed Color	Dark Gray
12	Cotyledon Color	Light red
13	Seed size	Large
14	Flower Color	Light Pink
15	Yield (Qt/ha)	Research Field On-farmer's field
		22-25 14-16
16	Disease reaction	Tolerant to rust, wilt and Aschochyta blight
17	Yield advantage over Asano (%)	29.76
18	Year of Release	2021
19	Breeder and Maintainer	SARC/ IQOO

Table 3. Mean grain yield (kg/ha) of 17 Lentil genotypes across locations and years.

Entry	Sinana			Agarfa			Mean	Yield Adv. over St. check
	2016	2017	2018	2016	2017	2018		
DZ -2012-LN-0051	1810	2404	3364	1690	1458	1623	2058	29.76%
DZ -2012-LN-0057	916	2018	3441	766	933	2113	1698	
DZ -2012-LN-0059	1363	2375	3308	614	694	861	1536	
DZ -2012-LN-00118	1484	2581	3089	1574	1412	1640	1963	
FLIP-96-49L	1384	2347	3278	1498	1272	1142	1820	
DZ -2012-LN-0038	1480	2360	3084	1991	1036	1338	1881	
DZ -2012-LN-00107	1408	2195	3300	2003	987	1175	1845	
DZ -2012-LN-0058	1220	1857	2853	1425	884	993	1539	
DZ -2012-LN-0048	1505	2542	3457	1195	916	1461	1846	
FLIP-97-33L	1194	2398	3021	1467	1138	1870	1848	
DZ -2012-LN-0065	1621	2329	3003	1173	611	1118	1643	
FLIP-86-38L	1635	2289	3614	1939	1135	1170	1964	
FLIP-89-19L	1284	1788	2323	1569	675	1885	1587	
DZ -2012-LN-0095	1715	2746	2678	1648	1019	2055	1977	
DZ -2012-LN-0085	1882	2836	3237	1838	1582	2496	2311	
Asano (St. check)	1211	1634	2020	1525	1032	2095	1586	
Local check	1805	1901	1345	1544	763	574	1322	
Means	1466	2271	3136	1498	1032	1506	1790	
LSD (<0.05)	479.5	590.0	907.5	565.2	634.6	716.1	291.9	
C.V	23.0	18.0	20.0	22.5	23.0	23.0	20.6	

Table 4. Mean Seed yield and other agronomic traits for 17 lentil genotypes tested in regional Variety Trial combined over two locations (Sinana and Agarfa) over three years (2016-2018).

Entry	DF	DM	Stand %	PH (cm)	NPP	NSP	HSW (g)	SY (kg/ha)
DZ -2012-LN-0051	62	122	76	32	33	1	3.7	2058
DZ -2012-LN-0057	64	126	77	34	31	1	4.0	1698
DZ -2012-LN-0059	63	125	75	33	37	1	3.1	1536
DZ -2012-LN-00118	61	122	78	31	33	1	3.6	1963
FLIP-96-49L	62	123	75	31	36	1	3.5	1820
DZ -2012-LN-0038	62	123	78	31	33	1	3.6	1881
DZ -2012-LN-00107	62	123	79	31	33	1	3.6	1845
DZ -2012-LN-0058	61	125	74	31	35	1	2.7	1539
DZ -2012-LN-0048	61	124	76	32	37	1	3.2	1846
FLIP-97-33L	62	123	76	32	35	1	3.4	1848
DZ -2012-LN-0065	61	125	77	35	36	1	3.0	1643
FLIP-86-38L	62	122	77	32	31	1	3.6	1964
FLIP-89-19L	61	125	79	32	30	1	3.6	1587
DZ -2012-LN-0095	63	125	78	34	39	1	2.7	1977
DZ -2012-LN-0085	62	125	79	34	37	2	2.8	2311
Asano (St. Check)	61	124	77	31	32	1	3.7	1586
Local check	62	122	74	33	33	1	2.4	1322
Mean	62	124	77	32	34	1	3.3	1790
LSD (<0.05)	0.8	1.2	3.3	2.0	7.7	0.2	0.1	291.9
CV%	2.4	1.7	7.5	10.8	23.9	22.6	7.1	20.6

Table-5. Mean seed yield, agronomic traits and disease reaction of 'Debine' along with standard and Local checks tested in two environments at varietal verification levels during 2015-2017 cropping seasons.

Entry	Agronomic traits								Disease Reaction (1-9)		
	DF	DM	Stand %	PH (cm)	NPP	NSP	HSW (g)	SY (kg/ha)	ASB	Rust	RR
DZ -2012-LN-0051	62	122	76	32	33	1	3.7	2058	4	3	3
Asano	61	124	77	31	32	1	3.7	1586	5	3	4
DZ -2012-LN-0085	62	125	79	34	37	2	2.8	2311	4	3	3
Local check	62	122	74	33	33	1	2.4	1322	5	5	4

Note: DF = days to 50% maturity, DM, days to 90% maturity, PH = plant height (cm), NPP = Number of pods per plant, NSP = Number of seed per plant, HSW = Hundred seed weight (g), GY = grain yield (kg), ASB = Aschocyta Blight, RR = Root Rot.

4. Conclusion

Grain yield is the primary trait of interest and a prime objective in Lentil breeding programs for many decades. "Debine" produced high yield, and it had a more stable performance in seed yield over locations and years than the standard check variety. The current variety, Debine has 29.76% yield advantages over the widely cultivated lentil varieties, Asano. Therefore, wide cultivation of Debine variety will boost productivity and marketability of the crop and improve farmers' income. Debine was resistant to major diseases of Lentil that prevailed in the growing areas. Farmers also preferred the variety for its superior performance over the existing local variety, which is manifested by good plant height, better pods load and number of branches per plant. Hence, Debine was verified and officially released for large scale production in major Lentil growing areas of Bale highland and other similar agro-ecologies.

Acknowledgements

The authors would like to thank Oromia Agricultural Research Institute for funding the research throughout the varietal development process. They also thank all staff members of the Pulse and Oil crops research case Team of

the Sinana Agricultural Research Center for their unreserved efforts in field trail management and data collection during the experimental period.

References

- [1] Aghili, Parisa. "Study of correlation and relationships between seed yield and yield components in Lentil (*Lens culinaris* Medik)." *Ann. Biol. Res* 3. 11 (2012): 5042-5045.
- [2] Ates, D., S. Aldemir, B. Yagmur, A. Kahraman, H. Ozkan, A. Vandenberg and M. B. Tanyolac. 2018a. QTL Mapping of Genome Regions Controlling Manganese Uptake in Lentil Seed. *G3: Genes, Genomes, Genetics* 8 (5): 1409-1416.
- [3] Ates, D., T. Sever, S. Aldemir, B. Yagmur, H. Y. Temel, H. B. Kaya, A. Alsaleh, A. Kahraman, H. Ozkan, A. Vandenberg and B. Tanyolac. 2016. Identification QTLs controlling genes for se uptake in lentil seeds. *PLoS One* 11: e0149210.
- [4] Central Statistical Agency (CSA), "Report on area and production of major crops (private peasant holdings, meher season)," *Statistical Bulletin*, vol. 590, pp. 19-21, 2020/21.
- [5] Frederick M, Cho S, Sarker A, McPhee K, Coyne C, et al. (2006) Application of biotechnology in breeding lentil for resistance to biotic and abiotic stress. *Euphytica* 147 (1-2): 149-165.

- [6] Iqbal A, Khalil IA, Ateeq N, Khan N (2006) Nutritional quality of important legumes. *Food chemistry* 97 (2): 331-335.
- [7] Muehlbauer FJ, Cubero JI, Summerfield RJ (1985) Lentil (*Lens culinaris* Medic.). In: Summerfield RJ & Roberts EH (Eds.), *Grain Legume Crops*. Collins, 8 Grafton Street, London, UK, pp. 266-311.
- [8] Sarker A, Kumar S (2011) Lentils in production and food systems in West Asia and Africa. International Center for Agricultural Research in the Dry Areas (ICARDA), Aleppo, Syria. *Grain Legumes* 57: 46-48.
- [9] Shrestha, R., Rizvi, A. H. Sarkar, A. Darai, R. Paneru, R. B. Vandenberg, A. and Singh, M. 2018. Genotypic Variability and Genotype x Environment Interaction for Iron and Zinc Content in Lentil under Nepalese Environments. Published in *Crop Sci.* 58: 2503–2510 (2018).
- [10] Toklu, F., T. Karakoy, E. Haklı, T. Bicer, A. Brandolini, B. Kilian and H. Ozkan. 2009. Genetic variation among lentil (*Lens culinaris* Medik) landraces from Southeast Turkey. *Plant Breeding* 128: 178-86.
- [11] Williams PC, Singh U (1988) Quality screening and evaluation in pulse breeding. In: Summerfield RJ (Ed.), *World Crops: Cool Season Food Legumes*. Kluwer Academic Publishers, Dordrecht, Netherlands, pp. 445-457.
- [12] Winch T (2006) *Growing food: a guide to food production*. Springer publisher, Netherlands, pp. 333. 13. Williams PC, Singh U (1988) Quality screening and evaluation in pulse breeding. In: Summerfield RJ (Ed.), *World Crops: Cool Season Food Legumes*. Kluwer Academic Publishers, Dordrecht, Netherlands, pp. 445-457.